Parameter	$\geq 32\%$ CL	half width	$\geq 5\%$ CL	half width
λ	0.2221 ± 0.0021		0.2221 ± 0.0041	
A	0.770 - 0.888	0.059	0.754 - 0.906	0.076
$\bar{\rho}$	0.08 - 0.35	0.14	0.04 - 0.38	0.17
$\bar{\eta}$	0.22 - 0.46	0.12	0.21 - 0.49	0.14
J (10^{-5})	2.0 - 3.5	0.8	1.9 - 3.7	0.9
$\sin 2\alpha$	-0.91 - 0.34	0.63	-0.96 - 0.49	0.73
$\sin 2\beta$	0.50 - 0.86	0.18	0.47 - 0.89	0.21
α	80° - 123°	22°	75° - 127°	26°
β	15.0° - 29.7°	7.4°	14.0° - 31.4°	8.7°
$\gamma = \delta$	37° - 75°	19°	34° - 82°	24°
$\sin\theta_{12}$	0.2221 ± 0.0021		0.2221 ± 0.0041	
$\sin\theta_{13}$ (10^{-3})	2.70 - 4.31	0.81	2.49 - 4.55	1.03
$\sin\theta_{23}$ (10^{-3})	38.4 - 43.2	2.4	37.9 - 43.6	2.8
$\lvert V_{ud}\rvert$	0.97504 ± 0.00049		0.97504 ± 0.00094	
$\lvert V_{us}\rvert$	0.2221 ± 0.0021		0.2221 ± 0.0042	
$\lvert V_{ub}\rvert$ (10^{-3})	2.70 - 4.31	0.81	2.49 - 4.55	1.03
$\lvert V_{cd}\rvert$	0.2220 ± 0.0021		0.2220 ± 0.0042	
$\lvert V_{cs}\rvert$	0.97422 ± 0.00056		0.97422 ± 0.00102	
$\lvert V_{cb}\rvert$ (10^{-3})	38.4 - 43.2	2.4	37.9 - 43.6	2.8
$\lvert V_{td}\rvert$ (10^{-3})	6.6 - 9.2	1.3	6.3 - 9.6	1.6
$\lvert V_{ts}\rvert$ (10^{-3})	37.7 - 42.8	2.6	37.3 - 43.2	3.0
$\lvert V_{tb}\rvert$	0.99907 - 0.99927	10×10^{-5}	0.99905 - 0.99929	12×10^{-5}
Δm_s (ps^{-1})	15.5 - 33.7	9.1	15.0 - 42.0	13.5
$\mathrm{BR}(K_\mathrm{L}^0 \to \pi^0\nu\bar{\nu})$ (10^{-11})	1.3 - 4.0	1.4	1.2 - 4.4	1.6
$\mathrm{BR}(K^+ \to \pi^+\nu\bar{\nu})$ (10^{-11})	5.1 - 9.6	2.3	4.8 - 10.5	2.9
$\mathrm{BR}(B^+ \to \tau^+\nu_\tau)$ (10^{-5})	4.6 - 20.0	7.7	3.6 - 23.6	10.0
$\mathrm{BR}(B^+ \to \mu^+\nu_\mu)$ (10^{-7})	1.8 - 7.9	3.1	1.5 - 9.3	3.9
$f_{B_d}\sqrt{B_d}$ (MeV)	193 - 271	39	184 - 284	50
B_K	> 0.55		> 0.50	
m_t (GeV)	106 - 406	150	93 - 565	236

Table 1: *Fit results for the various CKM parameters, the CKM matrix elements, branching ratios of some rare K and B meson decays and theoretical quantities. Ranges are given for the quantities that are limited by systematic theoretical errors.*